sequence that is at least 80% identical to any one of SEQ ID NOs: 1-7, 149-164, 333-337, and 384 and/or a V_L Comprising a sequence that is at least 80% identical to any one of SEQ ID NOs: 8-13, 148, 338-343, and 385. For example, the antibody, or fragment or variant thereof, may comprise: a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 6 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 12; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 7 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 13; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 333 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 338; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 334 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 339; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 335 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 340; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 336 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 341; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 337 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 342; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 335 and a V_I Comprising a sequence that is at least 90% identical to SEQ ID NO: 343; a V_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 384 and a V_z Comprising a sequence that is at least 90% identical to SEQ ID NO: 385; or a \bar{V}_H Comprising a sequence that is at least 90% identical to SEQ ID NO: 386 and a V_L Comprising a sequence that is at least 90% identical to SEQ ID NO: 387.

[0017] In certain embodiments, the anti-PD-1 antibody, or fragment or variant thereof, is: AM-0001; AMP-224; balstilimab; budigalimab; BI 754091; camrelizumab; cemiplimab; cetrelimab; dostarlinab; JTX-4014; MEDI-0680; MGA012; nivolumab; pembrolizumab; pidilizunab; prolgolimab; sasanlimab; sintilimab; spartalizumab; STI-1110; tislelizumab; toripalimab; or zimberelimab; or a fragment or variant thereof. For example, the anti-PD-1 antibody, or fragment or variant thereof, is: pembrolizumab; nivolumab; zimberelimab; or cetrelimab; or a fragment or variant thereof. In certain embodiments, the anti-PD-1 antibody, or fragment or variant thereof, of the first component is cetrelimab, or a fragment or variant thereof.

[0018] In certain embodiments, the anti-PD-1 antibody, or fragment or variant thereof, comprises a sequence that is at least 90% identical to SEQ ID NO: 15 and a sequence that is at least 90% identical to SEQ ID NO: 16 or 143. In certain embodiments, the anti-PD-1 antibody, or fragment or variant thereof, comprises a sequence that is at least 80% identical to SEQ ID NO: 296 and a sequence that is at least 80% identical to any one of SEQ ID NOs: 144, 145, and 295. In certain embodiments, the anti-PD-1 antibody, or fragment or variant thereof, comprises a sequence that is at least 80% identical to SEQ ID NO: 13 or 15 and a sequence that is at least 80% identical to SEQ ID NO: 294 or 295.

[0019] In certain embodiments, the second component of the fusion protein is a cytokine trap, for example a TGF- β cytokine trap. The TGF- β cytokine trap may, for example, comprise: a transforming growth factor receptor (TGF β R) or a fragment or variant thereof; an anti-TGF- β antibody or an antigen binding fragment or variant thereof; a TGF- β inhibi-

tory peptide or a fragment or variant thereof; and/or a $TGF-\beta$ antagonistic peptide or a fragment or variant thereof.

[0020] In certain embodiments, the TGF- β cytokine trap comprises a transforming growth factor beta receptor II (TGF β RII) or a fragment or variant thereof.

[0021] In certain embodiments, the TGF- β cytokine trap comprises a TGF β R extracellular domain or a fragment or variant thereof. In certain embodiments, the TGF- β cytokine trap comprises a TGF β RII extracellular domain, or a fragment or variant thereof, for example comprising a sequence that is at least 80% identical to any one of SEQ ID NOs: 14, 141, and 142. In certain embodiments, the TGF β RII extracellular domain, or a fragment or variant thereof, binds TGF- β 1 and/or TGF- β 3. In certain embodiments, the TGF β RII extracellular domain, or a fragment or variant thereof, binds TGF- β 1 and TGF- β 3 but does not bind TGF- β 2 or binds TGF- β 2 at a lower affinity than it does TGF- β 1 and TGF- β 3.

[0022] In certain embodiments, the TGF- β cytokine trap comprises a sequence that is at least 80% identical to any one of SEQ ID NOs: 14, 141, and 142.

[0023] In certain embodiments, the TGF- β cytokine trap comprises an anti-TGF- β antibody or an antigen binding fragment or variant thereof. The antibody, or antigen binding fragment or variant thereof, may for example comprise a variable region of a heavy chain (V_H) and a variable region of a light chain (V_L) . In certain embodiments, the antibody, or antigen binding fragment or variant thereof, may comprise a V_H comprising a sequence that is at least 80% identical to any one of SEQ ID NOs: 166, 168, 169, 171, 173, and 175 and/or a V_L comprising a sequence that is at least 80% identical to any one of SEQ ID NOs: 165, 167, 170, 172, 174, 176, and 178.

[0024] In certain embodiments, the TGF- β cytokine trap comprises a TGF- β inhibitory peptide or a fragment or variant thereof, for example comprising a sequence that is at least 80% identical to any one of SEQ ID NOs: 468-507 and 263-267. In certain embodiments, the TGF- β cytokine trap comprises two or more TGF- β inhibitory peptides, or fragments or variants thereof. The two or more TGF- β inhibitory peptides, or fragments or variants thereof, may be connected by linker(s).

[0025] In certain embodiments, the TGF- β cytokine trap comprises a TGF- β antagonistic peptide or a fragment or variant thereof. In certain embodiments, the TGF- β cytokine trap comprising two or more TGF- β antagonistic peptides, or a fragments or variants thereof. The two or more TGF- β antagonistic peptides, or fragments or variants thereof, may be connected by linker(s).

[0026] In certain embodiments, the second component of the fusion protein is an ADA or a fragment or variant thereof. [0027] In certain embodiments, the ADA, of fragment or variant thereof, is adenosine deaminase 2 (ADA2) or a fragment or variant thereof, for example one comprising a sequence that is at least 80% identical to any one of SEQ ID NOs: 273-279 and 284. In certain embodiments, the ADA2, or fragment or variant thereof, comprises at least one amino acid substitution or deletion.

[0028] In certain embodiments, the first and second components are connected by a linker. The linker may, for example, comprise: (G4S)_n, wherein n is 2, 3, 4, 5, or 6; (Gly)_n, wherein n is 6, 7, or 8; (EAAAK)_n, wherein n is 1, 2, 3, 4, 5, or 6; A(EAAAK)₄ALEA(EAAAK)₄A; and/or a sequence of any one of SEQ ID NOs: 17-34.